

Neutron-Proton High-Energy Charge Exchange Scattering

Y. Yan^{*} and R. Tegen[†]

School of Physics, Suranaree University of Technology,
Nakhon Ratchasima 30000, Thailand

T. Gutsche, V.E. Lyubovitskij and Amand Faessler
Institute for Theoretical Physics, Tübingen University,
Auf der Morgenstelle 14, D-72076 Tübingen, Germany

February 1, 2008

^{*}e-mail: yupeng@ccs.sut.ac.th

[†]Permanent Address: Department of Physics, University of the Witwatersrand, P.O. WITS
2050, Johannesburg, South Africa

Abstract

The high energy pn charge exchange scattering reaction is studied in an effective hadron model for the energy range $45.9 \text{ GeV}^2 \leq s \leq 414.61 \text{ GeV}^2$.

The main features of the observed differential cross section, the forward peak and the scaling behavior over a large energy region, are well reproduced.

PACS. Numbers: 13.75.Cs, 13.85.Dz, 14.20.Dh

Key Words: background contribution, meson-exchange

Nucleon-nucleon and nucleon-antinucleon scattering cross sections (total and differential) have been measured over a very large energy range (from threshold up to several TeV for the total center-of-mass energy). Such data are the major source of information on the hadronic and sub-hadronic properties of all matter[1]. In this communication we focus on an interesting sub-set of those data, namely the neutron-proton charge-exchange (CEX) differential cross sections $d\sigma/dt$ as a function of s which exhibit two main features: (i) a sharp forward peak which appears for $|t| < 0.02$ (GeV/c)², and (ii) a scaling behavior, the nearly form-invariant differential cross section depends on the lab momentum p in the form of $1/p^2$ over a wide energy region. These two features represent an enormous challenge for theoretical nucleon-nucleon models.

The generally quite successful relativistic meson-exchange models of the nucleon-nucleon interaction find it very difficult to explain (i) and (ii) without adhoc adjustments to the underlying model. In any of those meson-exchange models one would expect the lightest charged mesons (π and ρ) to dominate the *CEX* differential cross sections at least in the low- $|t|$ region. It turns out that the one-pion-exchange alone gives rise to the observed $1/p^2$ dependence of $d\sigma/dt$ for $|t| > 0.1$ GeV/c, but fails to explain $d\sigma/dt$ at $t = 0$ where this contribution is exactly zero.

It is well known since the 1960's that the rho-exchange does not improve the situation. In 1963 it was pointed out by Phillips[2] that the data are consistent with

a nearly constant background field. Following this original observation, a number of works attempting to explain the origin of such a background fall into two categories, namely Regge absorption models (for example, by Henyey *et al.*[3] and Kane *et al.*[4]) and Regge cut models (for example, by Chia[5]). Regge absorption models assume that the pion pole is modified by inelastic channels while Regge cut models evaluate double exchanges. Those models are with some success in reproducing the np CEX differential cross section.

More recently the np CEX scattering was studied by Gibbs and Loiseau[6] in a model of one- and two-pion exchanges between the quark constituents of the nucleon. It is noticed that in this model the one-pion exchange cannot give a reasonable result, especially for the forward peak. With the two-pion exchange included, the model fits the np CEX differential cross section quite well. However, only a very narrow energy region was considered. The P-wave part of the two-pion exchange is similar to the rho-exchange which leads to a forward peak. This part, however, is energy independent for small momentum transfers at high energy and, hence, cannot be expected to provide a major part of the missing contributions to the nucleon-proton CEX differential cross section over the observed range of energies.

The model - In this work we study np CEX scattering, based on that the interaction for the nucleon-nucleon elastic scattering is mainly mediated by the quark-antiquark sea which might be parametrized by mesons and a background.

We start with the Lagrangians of the $NN\pi$ and $NN\rho$ coupling

$$\mathcal{L}_{\pi NN} = g_{\pi NN} \bar{N} i \gamma^5 \vec{\pi} \vec{\tau} N \quad (1)$$

$$\mathcal{L}_{\rho NN} = g_{\rho NN} \bar{N} \gamma^\mu \vec{\rho}_\mu \vec{\tau} N + \frac{1}{4M_N} f_{\rho NN} \bar{N} \sigma^{\mu\nu} (\partial_\mu \vec{\rho}_\nu - \partial_\nu \vec{\rho}_\mu) \vec{\tau} N \quad (2)$$

with

$$N = \begin{pmatrix} p \\ n \end{pmatrix} \quad (3)$$

where p , n , $\vec{\pi}$ and $\vec{\rho}_\mu$ stand for proton, neutron, π and ρ fields, respectively. These Lagrangians are widely used in various literatures such as Ref.[7]. It is straightforward to calculate $d\sigma/dt$ for high energies (s is much larger than any mass scale involved) and low $|t|$ with the results

$$\left(\frac{d\sigma}{dt} \right)_\pi \sim t^2/s^2 \quad (4)$$

and

$$\left(\frac{d\sigma}{dt} \right)_\rho \sim \text{Constant} \quad (5)$$

for one π and one ρ exchanges, respectively. As discussed above, eq.(5) is inconsistent with the scaling behavior as stated in (ii), so the one-rho exchange cannot dominate the neutron-proton $(d\sigma/dt)_{CEX}$ at large s and low $|t|$. While giving the correct energy-dependence, eq.(4) fails to reproduce the the forward peak for $(d\sigma/dt)_{t \sim 0}$.

Obviously there is an additional contribution, beyond π - and ρ -exchange, necessary to explain the t - and s -dependence of $d\sigma/dt$.

We study the problem based on the argument that the nucleon-nucleon elastic scattering is mainly mediated by the quark-antiquark sea around the quark core of nucleons, and the quark-antiquark sea might be parameterised as a four-nucleon contact interaction and by various observed mesons.

We use a model Lagrangian $\mathcal{L}_{eff}$ which includes the free part for the nucleons $\mathcal{L}_N$ and the mesons $\mathcal{L}_M$, the meson-nucleon interaction $\mathcal{L}_{MNN}$ and the four-nucleon interaction term $\mathcal{L}_{4N}$ modelling the short range nucleon-nucleon interactions or the background:

$$\mathcal{L}_{eff}(x) = \mathcal{L}_N(x) + \mathcal{L}_M(x) + \mathcal{L}_{MNN}(x) + \mathcal{L}_{4N}(x), \quad (6)$$

where

$$\mathcal{L}_N(x) = \bar{N}(x)(i\gamma^\mu\partial_\mu - M_N)N(x), \quad (7)$$

$$\mathcal{L}_M(x) = -\frac{1}{2}\vec{\pi}(x)[\square + M_\pi^2]\vec{\pi}(x) + \frac{1}{2}\vec{\rho}_\mu(x)[g^{\mu\nu}(\square + M_\rho^2) - \partial^\mu\partial^\nu]\vec{\rho}_\nu(x), \quad (8)$$

$$\begin{aligned} \mathcal{L}_{MNN}(x) &= \int d^4y \bar{N}(x) i\gamma^5 \vec{\pi}(y) \vec{\tau} G_{\pi NN}(x-y) N(x) \\ &+ \int d^4y \bar{N}(x) \gamma^\mu \vec{\rho}_\mu(y) \vec{\tau} G_{\rho NN}(x-y) N(x) \\ &+ \frac{1}{4M_N} \int d^4y \bar{N}(x) \sigma^{\mu\nu} \vec{R}_{\mu\nu}(y) \vec{\tau} F_{\rho NN}(x-y) N(x), \end{aligned} \quad (9)$$

$$\mathcal{L}_{4N}(x) = -\frac{g^2}{M_N^2} \int d^4y \bar{N}(x) N(x) F_{4N}(x-y) \bar{N}(y) N(y). \quad (10)$$

$\vec{R}_{\mu\nu} = \partial_\nu \vec{\rho}_\mu - \partial_\mu \vec{\rho}_\nu$ is the strength of the ρ -meson field; $G_{\pi NN}$, $G_{\rho NN}$ and $F_{\rho NN}$ are form factors modelling the distribution of the meson cloud in nucleon [8]; F_{4N} is a form factor modelling the short-ranged nucleon-nucleon interaction or the background. The Fourier-transform of the vertex form factors is defined as:

$$G(F)_{MNN}(x) = \int d^4p e^{-ipx} G(F)_{MNN}(p^2) \quad (11)$$

$$F_{4N}(x) = \int d^4p e^{-ipx} F_{4N}(p^2) \quad (12)$$

The πNN coupling constant is fixed from the Goldberger-Treiman relation:

$$G_{\pi NN}(m_\pi^2) = g_A \frac{M_N}{F_\pi} \quad (13)$$

where $g_A = 1.267$, $F_\pi = 93$ MeV and $M_N = 938$ MeV are the experimental values of axial nucleon charge, weak pion decay constant and nucleon mass resulting in

$$\frac{G_{\pi NN}^2(m_\pi^2)}{4\pi} = 13.0 \quad (14)$$

For the ρNN coupling constants we use the values fixed in low and medium energy NN reactions [7]

$$\frac{G_{\rho NN}^2(m_\rho^2)}{4\pi} = 0.84, \quad \frac{F_{\rho NN}(m_\rho^2)}{G_{\rho NN}(m_\rho^2)} = 6.1. \quad (15)$$

The background field contribution, eq.(10), is reminiscent of the Nambo-Jona-Lasinio (NJL) 4-fermion interaction[9] which in the chain approximation (RPA) gives rise to a pionic mode and a scalar quark-antiquark condensate. Here we adjust the 4-fermion coupling strength g in eq.(10) to the experimental np data. Due to the analogy between eq.(10) and the NJL Lagrangian we will consider the background contribution to have similar properties as if a ' σ '-like object was exchanged between two nucleons, hence we will adopt a non-trivial vertex function.

The quark-antiquark substructure of N , π , and ρ is assumed to be manifest in the non-trivial meson-nucleon vertex functions. We study various forms of those vertex functions such as monopole, dipole, multipole and exponential forms. We find that the experimental data strongly suggest the monopole form for the πNN vertex function, and favor the tripole form for the ρNN vertex function. For the background contribution, we adopt a dipole form. The t -dependent vertex functions are defined as

$$F_{4N}(t) = \frac{1}{(1 - t/\Lambda^2)^2}, \quad (16)$$

$$G_{\pi NN}(t) = G_{\pi NN}(m_\pi^2) \frac{1 - m_\pi^2/\Lambda_\pi^2}{1 - t/\Lambda_\pi^2} \quad (17)$$

and

$$G(F)_{\rho NN}(t) = G(F)_{\rho NN}(m_\rho^2) = \left(\frac{1 - m_\rho^2/\Lambda_\rho^2}{1 - t/\Lambda_\rho^2} \right)^3. \quad (18)$$

Table 1: Coupling constants and cutoff parameters employed in Model A and B.

Models	$g_\pi^2/4\pi$	$g_\rho^2/4\pi$	f_ρ/g_ρ	$g^2/4\pi$	Λ [GeV]	Λ_π [GeV]	Λ_ρ [GeV]
Model A	13.0	0.84	6.1	5.5	0.26	0.52	–
Model B	13.0	0.84	6.1	5.0	0.24	0.50	1.0

Here, in addition to the coupling strength g , the cutoffs Λ , Λ_π and Λ_ρ are free parameters adjusted to the experimental data.

In Fig. 1 we indicate the theoretical results with the dashed lines for the one π exchange plus the background contribution (Model A) and the solid lines for the one π and one ρ exchanges plus the background contribution (Model B). All the relevant parameters are listed in Table 1 both for Model A and B. Here $g(f)_M \doteq G(F)_{MNN}(M^2)$ are input parameters while other parameters are adjusted to the observed differential cross sections of the pn charge exchange scattering. The energy range considered here is $45.9 \text{ GeV}^2 \leq s \leq 414.61 \text{ GeV}^2$. As apparent from Fig.1, the experimental data are well reproduced by Model B. Model A (one π exchange plus the background contribution) reproduces the data well up to $s = 109.68 \text{ GeV}^2$, and the ρ exchange contribution becomes significant when s is larger than 120 GeV^2 .

The np CEX differential cross section is well reproduced in a simple model of one π and one ρ exchanges including an additional background contribution. The

background contribution is important only for very small $|t|$. The one ρ exchange contribution is negligible below $s = 100 \text{ GeV}^2$, but becomes important for s larger than 200 GeV^2 , particularly for small $|t|$. The one π exchange is important for all energies, in particular for large $|t|$.

References

- [1] Review of Particle Physics, D.E. Groom *et al.*, Eur. Phys. J. C **15** (2000) 1;
Landolt-Börnstein, New Series, Group I, Vol.**7** (1973) and **9a** (1980)
- [2] R.J.N. Phillips, Phys. Lett. 4, 19 (1963); Nucl. Phys. B **2** (1967) 394.
- [3] F. Henyey, G.L. Kane, J. Pumplin, and M.H. Ross, Phys. Rev. **182** (1969) 1579.
- [4] G.L. Kane, F. Henyey, D.R. Richards, M.Ross, and G. Williamson, Phys. Rev. Lett. **25** (1970) 1519.
- [5] S.P. Chia, Phys. Rev. D **5** (1972) 2316; D **7** (1973) 1496; D **15** (1977) 290.
- [6] W.R. Gibbs, and B. Loiseau, Phys. Rev. C **50** (1994) 2742.
- [7] R. Machleidt, K. Holinde, and Ch. Elster, Phys. Rep. **149** (1987) 1.

[8] M.A. Ivanov, M.P. Locher, V.E. Lyubovitskij, Few-Body Syst. Few-Body Syst. **21** (1996) 131.

[9] V. Dmitrasinovic *et. al.*, Ann. Phys. (N.Y.) 238 (1995) 332; S.P. Klevansky, Rev. Mod. Phys. 64 (1992) 649.

Fig. 1 Theoretical predictions for $d\sigma^{CEX}/dt$ compared to experimental data (taken from the compilation of Ref. [1]). Dashed curve for Model A (one π -exchange plus background contribution); solid curve for Model B (one π and one ρ exchanges plus background contribution). Here s are in GeV^2 .




Figure 1

Figure 1 (continue)

